



Effect of nutrients on *in vitro* culture of *Morus alba* L. (White mulberry)

Ankit Pradhan, Vishwanathan AS, Basavaraja R.

Received: 3 April 2010 / Accepted: 12 April 2010 / Published online: 22 June 2010

Abstract

Morus alba L. is an economically important plant used extensively in sericulture, also possessing medicinal properties. *In vitro* culture provides an effective means to overcome the common problems encountered in conventional propagation by stem cuttings and seeds. Effect of phytohormones, growth adjuvants, sugars and essential minerals on the *in vitro* response of *M. alba* explants has been described in the study. Phytohormones and growth adjuvants resulted in explicit response in terms of organogenesis. Sucrose and glucose containing culture media elicited best response among sugars in nodal explants. Typical symptoms were exhibited by the growing shootlets when the medium was deprived of essential minerals such as nitrogen and sulphur.

Keywords: Phytohormones, Sugars, Minerals, Growth adjuvants.

Abbreviations

MS: Murashige and Skoog; 2,4-D: 2, 4- Dichlorophenoxy acetic acid; NAA: α - Naphthoxy Acetic Acid; Indole -3 - Butyric Acid; Kn: Kinetin; BAP: 6- Benzyl Amino Purine.

Vishwanathan AS.

Department of Biosciences, Sri Sathya Sai University,
Prasanthi Nilayam, Andhra Pradesh – 515 134 (India).

Web address:

<http://bioresonline.org/archives/A101.pdf>

Article citation:

Vishwanathan AS. 2010. Effect of nutrients on *in vitro* culture of *Morus alba* L. (White mulberry).
Bioresearch Bulletin 1: 008-012.



INTRODUCTION

The mulberry plant (*Morus alba* L.) is chiefly exploited by the sericulture industry for its foliage, used as feed for silk worm (*Bombyx mori* L.), and for feeding ruminants (Arabshahi-Delouee and Urooj, 2007). Depending on the location where it is grown, the plant is also valued for its fruit (consumed fresh, in juice or as a preserve) and aerial parts (stem and leaves are used as vegetable) (Venkatesh Kumar and Chauhan, 2008). *M. alba* also contains a considerable amount of dietary supplements such as proteins, carbohydrates, fats, fibers, essential minerals (Srivastava et al, 2006), ascorbic acid and β -carotene (Ercisli and Orhan, 2007).

The plant is known to possess anti-hyperglycemic and anti-hyperpigmentation activity (Lee et al, 2002). The root bark of the plant possesses astringent and anthelmintic properties, hypotensive activity, anti-tumour activity and antimicrobial activity and is used in treatment of cough and cold (Venkatesh Kumar and Chauhan, 2008). Traditional reports of the application of *M. alba* in treatment of atherosclerosis and diabetes were experimentally validated by Enkhmaa et al. (2005) and Shibata et al. (2007).

Stem cuttings and seeds have been used conventionally for propagation of mulberry plants which have assumed tremendous economic importance over the years. Genotype, environmental factors and physiological state of the cuttings play a significant role in determination of success of rooting (Lu, 2002). Propagation by seeds is undesirable owing to a long juvenile period (Vijaya Chitra and Padmaja, 2005) coupled with cross pollination which in turn leads to a high level of heterozygosity (Anis et al, 2003). *In vitro* culture or micropropagation is a sought-after alternative, providing an effective means of rapid propagation of uniform genotypes of the plant. Application of modern techniques for development of new high-yielding varieties and cultivars depends on the availability of an efficient *in vitro* regeneration system (Agarwal et al, 2004). This communication provides an account of studies on the *in vitro* response of stem-bit and nodal explants of *M. alba* to select phytohormones, growth adjuvants, minerals and sugars. This study was designed with the aim of identifying specific growth supplements which would have a positive influence on growth of *in vitro* plantlets. Another objective of the study was to evaluate the

occurrence of the known *in vivo* physiological effects of some of the selected supplements on the plantlets *in vitro*.

MATERIALS AND METHODS

Local varieties of *M. alba* were procured from local farms in and around Puttaparthi and grown in the experimental garden of the Sri Sathya Sai University, Prasanthi Nilayam campus. These constituted the mother plants for the explants used during the course of the *in vitro* study. The stem-bit and nodal explants were washed under running tap water for 30 minutes, surface sterilized using 0.1% (w/v) aqueous solution of Mercuric Chloride and rinsed five to six times with sterile double distilled water.

The surface-sterilized explants were inoculated on Murashige and Skoog medium (Murashige and Skoog, 1962) under a laminar flow hood (KLENZAIDS, India). The cultures were maintained at $24 \pm 2^\circ\text{C}$ under 16/8 h photoperiod at a light intensity of around 3000 lux.

MS medium fortified with auxins (2,4-D, NAA and IBA), cytokinins (Kn, BAP) and growth adjuvants (coconut milk and casein hydrolysate) as individual treatments was used in the study. MS medium without its constituent nitrogen and sulphate mineral stock solutions was used to study the effects on explant growth and development. Each of the phytohormones was used at a concentration of 2mg L^{-1} of MS medium and the growth adjuvants at 15% w/v or v/v. Aliquots of MS medium containing different sugars such as glucose, sucrose, fructose, mannose, maltose, lactose and galactose at a concentration of 3% (w/v) was used to study the effect of different sugars. Basal MS medium without addition of any supplements represented the control.

RESULTS AND DISCUSSION

Studies on the *in vitro* response of explants of *M. alba* have been undertaken to understand the effects of different nutrients with respect to organogenesis. This study focuses on the effect of phytohormones, growth adjuvants, sugars and minerals as observed in mulberry explants *in vitro*.

Nodal explants showed a better response as compared to stem-bits in the study of the effect of phytohormones on *in vitro* culture of *M. alba*. Callus initiation and proliferation occurred in MS medium supplemented with 2,4-D. The callus was semi-transparent and off-white in the beginning



gradually turning brownish in colour. Both NAA and IBA were effective in initiating a response in terms of shoot formation and growth; the former resulting in development of unbranched roots and the latter, profusely branched roots. The cytokinins Kn and BAP promoted shoot growth. Kn resulted in shoots with considerably longer internodal distance compared to any other treatment and BAP treatment led to formation of leaf clusters. Treatment with casein hydrolysate in MS medium inoculated with nodal explants gave good results in shoot formation while coconut milk was more effective in callogenesis. Coconut milk showed similar response using stem-bit explants.

Micropropagation studies on mulberry, most commonly, have used MS medium culture of nodal cuttings and axillary and apical buds (Thomas, 2002). Islam *et al.* (1993) observed that explants in medium containing 2,4-D showed best callogenesis and callus proliferation. Pattnaik and Chand (1997) and Bhau and Wakhlu (2003) obtained multiple shoots in BAP fortified medium which enhanced bud-break frequency with the addition of GA₃. The combination of NAA and BAP induced shoot proliferation in mulberry and NAA, in particular, was found to be essential for root induction by Ponchia and Gardiman (1991). The increased inter-nodal distance in shootlets obtained in Kn containing medium, in this study, has not been reported hitherto in the available literature.

Nodal explants cultured in MS medium containing glucose and sucrose responded by producing best results in shoot length, leaf growth and the development of a highly branched,

extensive root system. This was followed by medium containing fructose. Quite notably, medium containing galactose did not show any response. The results of the response of nodal explants to different sugars using MS medium are summarized in **Table 1**.

Sucrose is the principal product of carbon fixation in plants (Avigad and Dey, 1997). Other sugars have to be converted to glucose, the starting point of carbohydrate metabolism, or its derivatives to enter the glycolysis cycle (Brownleader *et al.*, 1997). The varied response to the different sugars reflects on the ability of the plant in converting these sugars into suitable glycolysis intermediates. Enomoto (1987) reported that sucrose containing media elicit the best response in *in vitro* mulberry culture. However Ohyama (1970), Ohyama and Oka (1976) and Vijaya Chitra and Padmaja (2002) report fructose as being more favourable. The present study revealed that sucrose is the best for morphogenic response in shoot length, leaf growth and size. These characteristics are highly advantageous for the sericulture industry.

Alteration in the mineral constitution of MS medium produced pronounced physiological effects that reflected in the morphology of the developing explants.

Table 2 outlines the different morphological changes that occurred in the response of stem-bit explants in the absence of nitrates and sulphates in MS medium.

Absence of nitrogen compounds in MS medium resulted in very poor morphogenic response of explants. Nitrogen supplied as ammonium (NH₄⁺) and nitrate (NO₃⁻) ions promotes

Table 1: Comparative effect of different sugars on morphogenic response of *M. alba* explants

Sugar used	Average shoot length (cm)	Average leaf width (cm)	Root system
Monosaccharides			
Glucose	5.7 ± 0.3	3.0 ± 0.1	Thick, long extensively branched
Fructose	4.6 ± 0.3	2.7 ± 0.2	Thin, short, few rootlets
Mannose	1.8 ± 0.1	1.8 ± 0.1	Thin, short, no rootlets
Disaccharides			
Sucrose	6.0 ± 0.4	3.9 ± 0.3	Thick, long extensively branched
Maltose	2.9 ± 0.1	1.8 ± 0.2	No response
Lactose	3.6 ± 0.1	2.6 ± 0.3	Thin, short, few rootlets

Table 2: Effect of minerals on explants of *M. alba*

Alteration in MS medium composition	Effect on growing explants
MS without Nitrogen (NH ₄ NO ₃ and KNO ₃)	Shoots got dried up; leaves were shriveled and gradually turned yellow; rooting was negligible
MS without Sulphates (MgSO ₄ , MnSO ₄ , FeSO ₄ and CuSO ₄)	Shoots were very short; leaves were small, drooping, light green to yellow in colour with black tips, roots were profusely branched

cation anion balance within the plant (Bloom, 1994). Under severe nitrogen deficiency, leaves turn yellow and eventually fall off (Taiz and Zeiger, 2002). Sulphur is a constituent of several co-enzymes and vitamins essential for plant metabolism. Its deficiency leads to downward hooking of leaves along with stunted growth of shoots (Taiz and Zeiger, 2002). Deficiency of sulphates also decreases chlorophyll content and rate of photosynthesis (Lopez et al, 1996).

CONCLUSION

The mulberry plant is reported to be the one of best plants for silkworm rearing (Thomas, 2002). The qualitative and quantitative improvements in this plant are very essential from the aspect of enhancement of the sericulture industry. Micropropagation is an established alternative to the conventional propagation methods for obtaining improved mulberry cultivars. The hormone treatments discussed herein provide means to obtain mulberry plants with desirable traits. Biochemical tests could be conducted on *in vitro* generated plants to ascertain changes in phytochemical constituents which may have a bearing on the feeding habits of silkworms reared on mulberry. Similar studies could be designed for obtaining stress resistant varieties of mulberry which would be of benefit to the silk industry.

ACKNOWLEDGEMENTS

The authors express their gratitude to the chancellor of Sri Sathya Sai University, Bhagawan Sri Sathya Sai Baba, for His constant guidance and support.

REFERENCES

- Agarwal S, Kanwar K and Sharma D. 2004.** Factors affecting secondary somatic embryogenesis and embryo maturation in morus alba l. *Sci Hort.* 102:359-368.
- Anis M, Faisal M and Singh S. 2003.** Micropropagation of mulberry (morus alba l.) through in vitro culture of shoot tip and nodal explants. *Plant Tissue Culture* 13:47-51.
- Arabshahi-Delouee S and Urooj A. 2007.** Antioxidant properties of various solvent extracts of mulberry (morus indica l.) leaves. *Food Chem.* 102:1233-1240.
- Avigad D and Dey P. 1997.** Carbohydrate metabolism: Storage carbohydrates. In: Dey P and Harborne J. *Plant biochemistry.* Academic Press, Great Britain.
- Bhau B and Wakhlu A. 2003.** Rapid micropropagation of five cultivars of mulberry. *Biol Plant.* 46:349-355.
- Bloom A. 1994.** Crop acquisition of ammonium and nitrate. In: Boote K, Bennett J, Sinclair T and Paulsen G. *Physiology and determination of crop yield.* Soil Science Society of America, Inc., Crop Science Society of America, Inc., Madison, WI.
- Brownleader M, Harborne J and Dey P. 1997.** Carbohydrate metabolism: Primary metabolism of monosaccharides In: Dey P and Harborne J. *Plant biochemistry.* Academic Press, Great Britain.
- Enkhmaa B, Shiwaku K, Katsube T, Kitajima K, Anuurad E, Yamasaki M and Yamane Y. 2005.** Mulberry (morus alba l.) leaves and their major



- flavonol quercetin 3-(6-malonylglucoside) attenuate atherosclerotic lesion development in ldl receptor-deficient mice. *J Nutr.* 135:729-734.
- Enomoto S. 1987.** Preservation of genetic resources of mulberry by means of tissue culture. *JARQ.* 21:205-210.
- Ercisli S and Orhan E. 2007.** Chemical composition of white (*morus alba*), red (*morus rubra*) and black (*morus nigra*) mulberry fruits. *Food Chem.*, 103:1380-1384.
- Islam R, Zaman A, Joarder O and Barman A. 1993.** In vitro propagation as an aid for cloning of *morus laevigata* wall. *Plant Cell Tiss Org Cult.* 33:339-341.
- Lee S, Choi S, Kim H, Hwang J, Lee B, Gao J and Kim S. 2002.** Mulberroside f isolated from the leaves of *morus alba* inhibits melanin biosynthesis. *Biol Pharm Bull.* 25:1045-1048.
- Lopez J, Tremblay N, Voogt W, Dubé S and Gosselin A. 1996.** Effects of varying sulphate concentrations on growth, physiology and yield of the greenhouse tomato. *Sci Hort.* 67:207-217.
- Lu M. 2002.** Micropropagation of *morus latifolia* poilet using axillary buds from mature trees. *Sci Hort.* 96:329-341.
- Murashige T and Skoog F. 1962.** A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol Plant.* 15:473-497.
- Ohyama K. 1970.** Tissue culture in mulberry tree. *JARQ.* 5:30-34.
- Ohyama K and Oka S. 1976.** Regeneration of whole plants from isolated shoot tips of mulberry tree. *The Journal of Sericulture Science of Japan.* 45:115-120.
- Pattnaik S and Chand P. 1997.** Rapid clonal propagation of three mulberries, *morus cathayana* hemsl, *m. Lhou koiz.* And *m. Serrata* roxb., through in vitro culture of apical shoot buds and nodal explants from mature trees. *Plant Cell Rep.*, 16:503-508.
- Ponchia G and Gardiman M. 1991.** Research on in vitro propagation of mulberry. *Acta Horticulturae.* 314:4-9.
- Shibata Y, Kume N, Arai H, Hayashida K, Inui-Hayashida A, Minami M, Mukai E, Toyohara M, Harauma A and Murayama T. 2007.** Mulberry leaf aqueous fractions inhibit tnf- α -induced nuclear factor $\kappa\beta$ (nf- $\kappa\beta$) activation and lectin-like oxidized ldl receptor-1 (lox-1) expression in vascular endothelial cells. *Atherosclerosis.* 193:20-27.
- Srivastava S, Kapoor R, Thathola A and Srivastava R. 2006.** Nutritional quality of leaves of some genotypes of mulberry (*morus alba*). *Int J Food Sci Nutr.* 57:305-313.
- Taiz L and Zeiger E. 2002.** Plant physiology Sinauer Associates, Sunderland.
- Thomas T. 2002.** Advances in mulberry tissue culture. *J Plant Biol.*, 45:7-21.
- Venkatesh Kumar R and Chauhan S. 2008.** Mulberry: Life enhancer. *J Med Pl Res.* 2:271-278.
- Vijaya Chitra D and Padmaja G. 2002.** Seasonal influence on axillary bud sprouting and micropropagation of elite cultivars of mulberry. *Sci Hort.* 92:55-68.
- Vijaya Chitra D and Padmaja G. 2005.** Shoot regeneration via direct organogenesis from in vitro derived leaves of mulberry using thidiazuron and 6-benzylaminopurine. *Sci Hort.*, 106:593-602.